	REPORT OF THE DEATH O	F A PATIENT TO THE CORONER
Surname: _	ROONEY	Christian Names: William
flex:		Marital State: Single
Admitted:	14.2.86. at 10 a.m.	Died: 20.2.86. at 2:20 3xMM p.m.
Next of Ki	n:	Relationship:
Address:		Telephone :
SYNOPS	IS OF CLINICAL NOTES (recorde	ed in narrative from to include):
pers inco	onnel under steps in all	istory): <u>Patient found by ambulance</u> Ley. Moving all limbs and speaking beying commands. Covered in blood. assault unknown.
or oth Youn limb maki fixe	er relevant clinical finding g man 25-35 years, B.P. s, arching back intermit	120/90, Heart Rate 120. Moving all tently, opening eyes spontaneously, as but not obeying commands. Pupils
Annual have the building that and an and the second s	etrical and brisk. Cont . No other injuries app	cusions over right anterior chest
c. Treatm	etrical and brisk. Cont . No other injuries app ment: <u>Narcan 0.4 mg. x 2</u> esponse. Skull x-ray -	usions over right anterior chest parent. 2 I.V.; 50% Dextrose 50 ml. I.V multiple skull fractures around
c. Treatm no r vaul I.V. Teta	etrical and brisk. Cont . No other injuries app nent: Narcan 0.4 mg. x 2 esponse. Skull x-ray - t and base. Mannitol 20 Flucloxacillin 1 g. q0 nus Toxoid. Dilantin 10	usions over right anterior chest barent. 2 I.V.; 50% Dextrose 50 ml. I.V multiple skull fractures around 0% 100 ml. I.V. Dexamethasone 12 mg. bh, I.V., Penicillin 600 mg. q6h I.V 00 mg. I.V. Endotracheal tube
c. Treatm no r vaul I.V. Teta inse Fole	etrical and brisk. Cont . No other injuries app ment: Narcan 0.4 mg. x 2 esponse. Skull x-ray - t and base. Mannitol 20 Flucloxacillin 1 g. q6 nus Toxoid. Dilantin 10 rted. Paralysed and ver y cather inserted. Trar	usions over right anterior chest barent. 2 I.V.; 50% Dextrose 50 ml. I.V multiple skull fractures around 0% 100 ml. I.V. Dexamethasone 12 mg. 5h, I.V., Penicillin 600 mg. g6h I.V
c. Treatm no r vaul I.V. Teta inse Fole d. Subseq Unab	etrical and brisk. Cont . No other injuries app nent: Narcan 0.4 mg. x 2 esponse. Skull x-ray - t and base. Mannitol 20 Flucloxacillin 1 g. q6 nus Toxoid. Dilantin 10 rted. Paralysed and ver y cather inserted. Tran puent progress: Care	2 I.V.; 50% Dextrose 50 ml. I.V multiple skull fractures around 0% 100 ml. I.V. Dexamethasone 12 mg. 6h, I.V., Penicillin 600 mg. q6h I.V 100 mg. I.V. Endotracheal tube ttilated. Nasogastric tube inserted sferred to C.T. Scan with Intensive a Escort.
c. Treatm no r vaul I.V. Teta inse Fole d. Subseq Unab from	etrical and brisk. Cont . No other injuries app ment: Narcan 0.4 mg. x 2 esponse. Skull x-ray - t and base. Mannitol 20 Flucloxacillin 1 g. q6 nus Toxoid. Dilantin 10 rted. Paralysed and ver y cather inserted. Tran uent progress: Care le to comment as not inv Accident and Emergency.	2 I.V.; 50% Dextrose 50 ml. I.V multiple skull fractures around 0% 100 ml. I.V. Dexamethasone 12 mg. 6h, I.V., Penicillin 600 mg. q6h I.V 100 mg. I.V. Endotracheal tube ttilated. Nasogastric tube inserted sferred to C.T. Scan with Intensive a Escort.
c. Treatm no r vaul I.V. Teta inse Fole d. Subseq Unab from	etrical and brisk. Cont . No other injuries app ment: Narcan 0.4 mg. x 2 esponse. Skull x-ray - t and base. Mannitol 20 Flucloxacillin 1 g. q6 nus Toxoid. Dilantin 10 rted. Paralysed and ver y cather inserted. Tran uent progress: Care le to comment as not inv Accident and Emergency.	2 I.V.; 50% Dextrose 50 ml. I.V multiple skull fractures around 0% 100 ml. I.V. Dexamethasone 12 mg. 0h, I.V., Penicillin 600 mg. q6h I.V 00 mg. I.V. Endotracheal tube ttilated. Nasogastric tube inserted asferred to C.T. Scan with Intensive e Escort. volved with patient after transfer
c. Treatm no r vaul I.V. Teta inse Fole d. Subseq Unab from	etrical and brisk. Cont . No other injuries app ment: Narcan 0.4 mg. x 2 esponse. Skull x-ray - t and base. Mannitol 20 Flucloxacillin 1 g. q6 nus Toxoid. Dilantin 10 rted. Paralysed and ver y cather inserted. Tran uent progress: Care le to comment as not inv Accident and Emergency.	2 I.V.; 50% Dextrose 50 ml. I.V multiple skull fractures around 0% 100 ml. I.V. Dexamethasone 12 mg. 0h, I.V., Penicillin 600 mg. q6h I.V 00 mg. I.V. Endotracheal tube ttilated. Nasogastric tube inserted asferred to C.T. Scan with Intensive e Escort. volved with patient after transfer
c. Treatm no r vaul I.V. Teta inse fole d. Subseq unab from	etrical and brisk. Cont . No other injuries app ment: Narcan 0.4 mg. x 2 esponse. Skull x-ray - t and base. Mannitol 20 Flucloxacillin 1 g. q6 nus Toxoid. Dilantin 10 rted. Paralysed and ver y cather inserted. Tran uent progress: Care le to comment as not inv Accident and Emergency.	2 I.V.; 50% Dextrose 50 ml. I.V multiple skull fractures around 0% 100 ml. I.V. Dexamethasone 12 mg. 0h, I.V., Penicillin 600 mg. q6h I.V 00 mg. I.V. Endotracheal tube ttilated. Nasogastric tube inserted asferred to C.T. Scan with Intensive e Escort. volved with patient after transfer

1
